

Precision measurement of the lifetime of the $6p\ ^2P_{1/2}$ level of Yb^+

S. Olmschenk, D. Hayes, D. N. Matsukevich, P. Maunz, and C. Monroe
JQI and Department of Physics, University of Maryland, College Park, Maryland, 20742, USA

We present a precise measurement of the lifetime of the $6p\ ^2P_{1/2}$ excited state of a single trapped ytterbium ion (Yb^+). We use a time-correlated single photon-counting technique [1] adapted to utilize the features of a single-atom system [2]. In particular, ultrafast pulses excite a single trapped Yb^+ ion and the emitted photons are coupled into a single-mode optical fiber. By performing the measurement on a single atom with fast excitation and excellent spatial filtering, we are able to eliminate common systematics. Among other things, experimental measurements of Yb-like ions may be used to test *ab initio* atomic structure calculations [3].

This work is supported by IARPA under ARO contract, the NSF PIF Program, and the NSF Physics Frontier Center at JQI.

-
- [1] L. Young, W. T. Hill III, S. J. Sibener, S. D. Price, C. E. Tanner, C. E. Wieman, and S. R. Leone, *Phys. Rev. A* **50**, 2174 (1994).
 - [2] D. L. Moehring, B. B. Blinov, D. W. Gidley, R. N. Kohn, Jr., M. J. Madsen, T. D. Sanderson, R. S. Vallery, and C. Monroe, *Phys. Rev. A* **73**, 023413 (2006).
 - [3] U. I. Safronova, W. R. Johnson, M. S. Safronova, and J. R. Albritton, *Phys. Rev. A* **66**, 022507 (2002).